

The objective of this test is to test whether the null hypothesis represents 'non-inferiority' or 'superiority' depends on the context and whether the non-inferiority/superiority margin,  $\delta$ , is positive or negative. In this setting, we wish to test whether a mean,  $\mu$ , is non-inferior/superior to a reference value,  $\mu_0$ . The idea is that statistically significant differences between the mean and the reference value may not be of interest unless the difference is greater than a threshold,  $\delta$ . This is particularly popular in clinical studies, where the margin is chosen based on clinical judgement and subject-domain knowledge. The hypotheses to test are

$$\begin{aligned} H_0 &: \mu - \mu_0 \leq \delta \\ H_1 &: \mu - \mu_0 > \delta \end{aligned}$$

and  $\delta$  is the superiority or non-inferiority margin.

## Formulas

This calculator uses the following formulas to compute sample size and power, respectively:

$$n = \left( \sigma \frac{z_{1-\alpha} + z_{1-\beta}}{\mu - \mu_0 - \delta} \right)^2$$

$$1 - \beta = \Phi(z - z_{1-\alpha}) + \Phi(-z - z_{1-\alpha}) \quad , \quad z = \frac{\mu - \mu_0 - \delta}{\sigma/\sqrt{n}}$$

where

$n$  is sample size

$\sigma$  is standard deviation

$\Phi$  is the [standard Normal distribution function](#)

$\Phi^{-1}$  is the [standard Normal quantile function](#)

$\alpha$  is Type I error

$\beta$  is Type II error, meaning  $1 - \beta$  is power

$\delta$  is the testing margin

## References

Chow S, Shao J, Wang H. 2008. Sample Size Calculations in Clinical Research. 2nd Ed. Chapman & Hall/CRC Biostatistics Series. page 52.